



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/00	A2	(11) International Publication Number: WO 99/40904 (43) International Publication Date: 19 August 1999 (19.08.99)
(21) International Application Number: PCT/EP99/00614 (22) International Filing Date: 30 January 1999 (30.01.99) (30) Priority Data: 98/01709 12 February 1998 (12.02.98) FR (71) Applicant (for all designated States except US): MERCK PATENT GMBH [DE/DE]; Frankfurter Strasse 250, D-64293 Darmstadt (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): BONHOMME, Yves [FR/FR]; 21 avenue de la Paix, F-69260 Charbonnières les Bains (FR). BRIET, Philippe [FR/FR]; 216, avenue Félix Faure, F-69003 Lyon (FR). (74) Common Representative: MERCK PATENT GMBH; Postfach, D-64271 Darmstadt (DE).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: PHARMACEUTICAL COMPOSITION COMPRISING A COMBINATION OF METFORMIN AND FIBRATE, AND ITS USE FOR THE PREPARATION OF MEDICINES INTENDED TO REDUCE HYPERGLYCAEMIA (57) Abstract The present invention relates to a pharmaceutical composition comprising, as active principles, (i) metformin optionally in the form of one of its pharmaceutically acceptable salts, and (ii) a fibrate chosen from fenofibrate and bezafibrate, in combination with one or more pharmaceutically acceptable excipients. These compositions are particularly suitable for the treatment of non-insulin-dependent diabetes.		

BEST AVAILABLE COPY

BEST AVAILABLE COPY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakistan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

Pharmaceutical composition comprising a combination of metformin and fibrate, and its use for the preparation of medicines intended to reduce hyperglycaemia

- 1 -

The invention relates to a pharmaceutical composition containing a combination of metformin and of a fibrate chosen from fenofibrate and bezafibrate, as active principles. The invention also relates to the
5 use of metformin and of a fibrate chosen from fenofibrate and bezafibrate for the preparation of a medicinal combination intended to reduce the hyperglycaemia of non-insulin-dependent diabetes.

Metformin is mainly known for its anti-
10 hyperglycaemic activity and is widely used in the treatment of non-insulin-dependent diabetes. In the case of insulin-dependent diabetes, metformin is also administered to the patient in combination with insulin.

Bezafibrate and fenofibrate belong to the family of fibrates whose anti-hyperlipidic properties are well known. More specifically, the fibrates act on hypercholesterolaemia and hypertriglyceridaemia by inducing a reduction in the total cholesterol level as
20 well as the cholesterol linked to low density lipoproteins (LDL-cholesterol) and an even greater reduction in the levels of triglycerides and in particular of triglycerides linked to very low density lipoproteins (VLDL-triglycerides).

Bezafibrate has already been administered to non-insulin-dependent diabetics on account of its hypolipaeamic properties. This is because non-insulin-dependent diabetes is often accompanied by serious lipid metabolism disorders; consequently, one of the
30 main causes of mortality of patients suffering from this type of diabetes is the appearance of coronary diseases or disorders of the cerebrovascular system or of the peripheral vascular system which can lead to myocardial infarction.

The value of a treatment with bezafibrate in the case of diabetics suffering from non-insulin-dependent diabetes has been reported in particular by P. W. Seviour et al. in Diabetic Medicine, Vol. 5, 166-171 (1988).

The combination of a hypoglycaemic agent and of an anti-lipaemic agent has already been envisaged in the art, and especially for treating diabetics also displaying hyperlipaemia. Contradictory results were
5 obtained depending on the nature of the active substances. The study by A. K. Jain et al. published in Diabetes, Vol. 34, 1985, Vol. 293 (25), 1283 shows, for example, that better control of the hyperglycaemia is obtained by joint administration of sulphonylurea
10 (hypoglycaemic agent) and of halogenate (antilepaemic agent). However, that document reveals the absence of an effect of clofibrate (a known antilipaemic agent) on the seric glucose level in diabetic patients treated simultaneously with sulphonylurea.

15 Among the studies relating to combined therapies, mention may also be made of the combination of metformin and clofibrate proposed by S. R. De Silva et al. in Diabete & Metabolisme, 1979, 5, 223-229. That author notes a slight improvement in the hypoglycaemia
20 on simultaneous administration of clofibrate and metformin. However, the essential advantage of this combination lies manifestly in the parallel reduction of the levels of cholesterol and of triglycerides. It thus results from that publication that the overall
25 effect of the combination is the simple addition of the respective effects of each of the active substances.

Surprisingly, the present inventors have discovered that a specific combination of a
30 hypoglycaemic agent with an antilipaemic agent leads to a significant improvement of the hyperglycaemia in a diabetic patient suffering from non-insulin-dependent diabetes. More specifically, a synergistic effect has been obtained by combined administration of metformin and of a fibrate chosen from fenofibrate and
35 bezafibrate. The same advantageous results have been observed using a pharmaceutically acceptable salt of metformin in combination with one of these two fibrates.

The synergistic effect observed lies in a marked improvement of the hypoglycaemia, this being found both in patients with hyperlipaemia and in non-dyslipidaemic patients.

5 Thus, the invention relates to a pharmaceutical composition comprising, as active principles, (i) metformin optionally in the form of one of its pharmaceutically acceptable salts, and (ii) a fibrate
10 chosen from fenofibrate and bezafibrate, in combination with one or more pharmaceutically acceptable excipients.

15 This composition is more particularly suitable for reducing the hyperglycaemia of non-insulin-dependent diabetes. It can also be used on non-dyslipidaemic patients.

20 According to the invention, the metformin can be administered in the form of one of its pharmaceutically acceptable salts, such as the hydrochloride, acetate, benzoate, citrate, fumarate, embonate, chlorophenoxyacetate, glycolate, palmoate, aspartate, methanesulphonate, maleate, parachlorophenoxyisobutyrate, formate, lactate, succinate, sulphate, tartrate, cyclohexanecarboxylate, hexanoate, octonate, decanoate, hexadecanoate,
25 octodecanoate, benzenesulphonate, trimethoxybenzoate, paratoluenesulphonate, adamantanecarboxylate, glycoxylate, glutamate, pyrrolidonecarboxylate, naphthalenesulphonate, 1-glucosephosphate, nitrate, sulphite, dithionate or phosphate.

30 Among these salts, the hydrochloride, fumarate, embonate and chlorophenoxyacetate are more particularly preferred.

35 The pharmaceutically acceptable salts of metformin are obtained in a manner which is known per se by the action of metformin on the corresponding acid.

 The compositions of the invention contain therapeutically effective amounts of the various active

principles. The ratios of the respective amounts of metformin and of fibrate thus vary in consequence.

Preferably, the weight ratio of metformin or of its pharmaceutically acceptable salt to fibrate ranges from 1:1 to 20:1, preferably from 1:1 to 5:1 and better still from 2:1 to 5:1.

The compositions of the invention are preferably administered parenterally, or better still orally, although the other routes of administration, for instance such as rectal administration, are not excluded.

When oral administration is envisaged, the compositions of the invention are in the form of gelatin capsules, effervescence tablets, coated or uncoated tablets, sachets, sugar-coated tablets, drinkable vials or solutions, microgranules or sustained-release forms.

When parenteral administration is envisaged, the compositions of the invention are in the form of injectable solutions and suspensions packaged in vials or bottles for slow venous infusion.

The forms for oral administration are prepared by mixing the active substance with various types of excipients or of vehicles, such as fillers, disintegration (or crumbling) agents, binders, dyes, flavour enhancers and the like, followed by shaping the mixture.

The dye can be any dye authorized for pharmaceutical use.

Examples of flavour enhancers include cocoa powder, mint, borneol and cinnamon powder.

Examples of binders which may be mentioned are polyvinylpyrrolidone, hydroxypropylmethylcellulose, alginic acid, carbomer, carboxymethylcellulose, dextrin, ethylcellulose, starch, sodium alginate, polymethacrylate, maltodextrin, liquid glucose, magnesium aluminium silicate, hydroxyethylcellulose, ethylcellulose, methylcellulose and guar gum.

It is possible to use alginic acid, sodium carboxymethylcellulose, colloidal silicon dioxide, sodium croscarmellose, crospovidone, guar gum, magnesium aluminium silicate, methylcellulose, microcrystalline cellulose, potassium polacrilin, cellulose powder, pregelatinized starch, sodium alginate or sodium starch glycolate as disintegration agent.

The fillers are, for example, cellulose, lactose, calcium hydrogenophosphate and microcrystalline cellulose.

The tablets can be obtained in a conventional manner by compressing granules in the presence of one or more lubricants. Suitable lubricants are calcium stearate, glyceryl monostearate, glyceryl palmitostearate, hydrogenated castor oil, hydrogenated plant oil, light mineral oil, magnesium stearate, polyethylene glycol, sodium benzoate, sodium lauryl sulphate, stearyl sodium fumarate, stearic acid, talc and zinc stearate. These tablets can then be coated using polymers in solution or suspension, such as hydroxypropylmethylcellulose or ethylcellulose.

The granules used to do this are prepared, for example, by using the wet-route granulation process starting with a mixture of the active principles with one or more excipients such as a binder, a crumbling agent (or disintegration agent) and a filler.

To obtain hard capsules, the mixture of active principles with a suitable filler (for example lactose) is incorporated into empty gelatin capsules optionally in the presence of a lubricant such as magnesium stearate, stearic acid, talc or zinc stearate.

Soft gelatin capsules are prepared by dissolving the active principles in a suitable solvent (for example polyethylene glycol), followed by incorporation into soft capsules.

The forms for parenteral administration are obtained in a conventional manner by mixing the active principles with buffers, stabilizers, preserving

agents, solubilizing agents, tonicity agents and suspension agents. In accordance with the known techniques, these mixtures are subsequently sterilized and then packaged in the form of intravenous
5 injections.

As buffer, a person skilled in the art can use buffers based on salts of organic phosphate.

Examples of suspension agents include methylcellulose, hydroxyethylcellulose, acacia and
10 sodium carboxymethylcellulose.

Examples of solubilizing agents include castor oil solidified with polyoxyethylene, polysorbate 80, nicotinamide or macrogol.

In addition, stabilizers which are useful
15 according to the invention are sodium sulphite and sodium metasulphite, while mention may be made of sodium p-hydroxybenzoate, sorbic acid, cresol and chlorocresol as preserving agents. For the preparation of an oral solution or suspension, the active
20 principles are dissolved or suspended in a suitable vehicle with a dispersing agent, a wetting agent, a suspension agent (for example polyvinylpyrrolidone), a preserving agent (such as methylparaben or propylparaben), a flavour enhancer or a dye.

25 For the preparation of suppositories, the active principles are mixed in a manner which is known per se with a suitable base constituent, such as polyethylene glycol or semi-synthetic glycerides.

For the preparation of microcapsules, the
30 active principles are combined with suitable diluents, suitable stabilizers, agents which promote sustained release of the active substances or any other type of additive for the formation of a central core which is then coated with a suitable polymer (for example a
35 water-soluble resin or a water-insoluble resin). The techniques known to those skilled in the art will be used for this purpose.

The microcapsules thus obtained are then optionally formulated in suitable dosage units.

A subject of the present invention is also the use of metformin optionally in the form of one of its pharmaceutically acceptable salts in combination with a fibrate chosen from bezafibrate and fenofibrate, for the preparation of a medicinal combination intended to reduce the hyperglycaemia of non-insulin-dependent diabetes.

According to another of its aspects, the invention relates to the use of metformin optionally in the form of one of its pharmaceutically acceptable salts, in combination with the said fibrate, for the preparation of a medicinal combination intended to reduce the hyperglycaemia of non-insulin-dependent diabetes in non-dyslipidaemic patients.

According to the invention, the term "medicinal combination" is intended to refer either to a pharmaceutical composition as defined above, in which the two active principles are the essential constituents of the same composition, or to a kit comprising two separate compositions, the first comprising metformin or its pharmaceutically acceptable salt as sole active principle, and the second comprising fibrate as sole active principle.

When the medicinal combination is in the form of a kit, the administration of the two compositions constituting this kit, although carried out separately, is simultaneous for a combined treatment.

The metformin can be in the form of any one of the salts defined above; however, it is preferred to use metformin as it is or in the form of the hydrochloride, fumarate, embonate or chlorophenoxyacetate.

According to a preferred embodiment, the amount of metformin or of its salt which is used is from one to twenty times the mass of the fibrate, preferably from one to five times and better still from two to five times.

When the metformin or its salt and the fibrate are incorporated into the same unit dose, the unit dose preferably comprises from 100 to 1000 mg of metformin.

In this case, the unit dose advantageously
5 comprises from 50 to 300 mg of fenofibrate or from 50 to 600 mg of bezafibrate.

The dosage naturally depends on the mode of administration, the therapeutic indication and the patient's age and condition.

10 In general, the daily dosage ranges between 100 and 2000 mg of metformin, between 50 and 600 mg of fenofibrate and between 50 and 1200 mg of bezafibrate.

The use of the compositions of the invention and the advantage of the use claimed are illustrated
15 hereinbelow with reference to the example which follows.

EXAMPLE

The synergism of action was proven using an
20 animal model. Non-insulin-dependent diabetes (NIDD) is induced by injecting streptozotocin into male Wistar rats. The action of clofibrate alone, of bezafibrate alone, of fenofibrate alone and of metformin alone was first evaluated in terms of glycaemia, cholesterol
25 level and level of triglycerides. Next, the metformin + clofibrate, metformin + bezafibrate and metformin + fenofibrate combinations were studied.

The procedure followed is as follows.

45 mg/kg of streptozotocin (STZ) dissolved in
30 physiological saline are administered to male Wistar rats. Two weeks after this treatment, blood is taken and the glycaemia is measured. Only the animals with a glycaemia of between 2 grams and 3 grams per litre are used for the treatments (about 6/10). The animals then
35 receive, orally, either metformin alone, or a fibrate alone, or a combination of the two in the doses indicated in Table 1 below. 23 days after the injection of streptozotocin, the animals are sacrificed and the following parameters are determined: glycaemia,

cholesterol and triglycerides. The averages obtained from 10 rats per group are modified by the standard error of the mean. The Student t test is carried out to evaluate the significance of the results obtained.

5 The combined results are reported in Table 1 below:

TABLE 1

Treatment	Glycaemia g/l	Cholesterol g/l	Triglycerides g/l
Absolute control	1.06 ± 0.06	0.50 ± 0.02	0.86 ± 0.04
Streptozotocin alone	2.68 ± 0.06 ^{oo}	0.65 ± 0.04 ^{oo}	1.25 ± 0.07 ^{oo}
Metformin (50 mg/kg)	1.74 ± 0.14 ^{**}	0.61 ± 0.03	0.85 ± 0.10 ^{**}
Clofibrate (100 mg/kg)	2.58 ± 0.11	0.63 ± 0.03	0.93 ± 0.11 [*]
Fenofibrate (50 mg/kg)	1.92 ± 0.20 ^{**}	0.45 ± 0.02 ^{**}	0.61 ± 0.10 ^{**}
Bezafibrate (50 mg/kg)	2.20 ± 0.21 [*]	0.52 ± 0.05 ^{**}	0.81 ± 0.11 ^{**}
Clofibrate (100 mg/kg) + Metformin (50 mg/kg)	1.72 ± 0.09 ^{**}	0.63 ± 0.02	0.94 ± 0.08 ^{**}
Fenofibrate (50 mg/kg) + Metformin (50 mg/kg)	1.44 ± 0.11 ^{**}	0.56 ± 0.05	0.63 ± 0.12 ^{**}
Bezafibrate (50 mg/kg) + Metformin (50 mg/kg)	1.43 ± 0.05 ^{**§}	0.48 ± 0.04 ^{**§}	0.62 ± 0.02 ^{**§}
^o p > 0.01 Comparison between absolute controls and NIDD STZ rats [*] p > 0.05, ^{**} p > 0.01 Comparison between control and treated STZ [§] p > 0.05 Comparison between metformin alone and in combination			

10

Examination of the results obtained quite clearly shows the synergism of action of the (metformin + fenofibrate) or (metformin + bezafibrate) combination on glycaemia. Whereas metformin alone leads to a glycaemia of 1.74 g/l and bezafibrate alone leads to a glycaemia of 2.20 g/l and fenofibrate alone leads to a glycaemia of 1.92 g/l, the metformin + bezafibrate/fenofibrate combinations lead, respectively, to glycaemias of 1.43 g/l and 1.44 g/l.

15

On the other hand, no synergism is observed for the clofibrate + metformin combination; in fact, the resulting glycaemia of 1.72 g/l is virtually that resulting from the administration of clofibrate.

5 Interestingly, it is more noted that:

- the fibrates, in particular bezafibrate and fenofibrate, when administered alone, show evidence of antihyperglycaemic properties. This effect may be associated with the enzymatic inductive effects by the action on glucose-6-phosphatase (action correlated with the variation of antipyrine);

10 - metformin, administered alone, possesses, besides its antidiabetic activity, effects on decreasing the levels of cholesterol and of triglycerides (in animals, such as in man).

15 This example unequivocally illustrates the surprising effect observed during the simultaneous administration of metformin and of a fibrate chosen from fenofibrate and bezafibrate.

CLAIMS

1. Pharmaceutical composition comprising, as active principles, (i) metformin optionally in the form of one of its pharmaceutically acceptable salts, and (ii) a fibrate chosen from fenofibrate and bezafibrate, in combination with one or more pharmaceutically acceptable excipients.
2. Composition according to Claim 1, for reducing the hyperglycaemia of non-insulin-dependent diabetes.
3. Composition according to either of Claims 1 and 2, for reducing the hyperglycaemia of non-insulin-dependent diabetes in non-dyslipidaemic patients.
4. Pharmaceutical composition according to any one of Claims 1 to 3, characterized in that the weight ratio of metformin or of its pharmaceutically acceptable salt to fibrate ranges from 1:1 to 20:1, preferably from 2:1 to 5:1.
5. Pharmaceutical composition according to any one of the preceding claims, characterized in that the metformin salt is a hydrochloride, fumarate, embonate or chlorophenoxyacetate.
6. Composition according to any one of the preceding claims, which is suitable for oral administration.
7. Use of metformin, optionally in the form of one of its pharmaceutically acceptable salts, in combination with a fibrate chosen from bezafibrate and fenofibrate, for the preparation of a medicinal combination intended to reduce the hyperglycaemia of non-insulin-dependent diabetes.
8. Use according to Claim 7, for the preparation of a medicinal combination intended to reduce the hyperglycaemia of non-insulin-dependent diabetes in non-dyslipidaemic patients.
9. Use according to either of Claims 7 and 8, characterized in that the metformin salt is a hydrochloride, a fumarate, an embonate or a chlorophenoxyacetate.

10. Use according to any one of the preceding claims, characterized in that the medicinal combination is in the form of a unit dose containing metformin, or one of its pharmaceutically acceptable salts, and the
5 fibrate.
11. Use according to Claim 10, characterized in that the unit dose comprises from 100 to 1000 mg of metformin and from 50 to 300 mg of fenofibrate.
12. Use according to Claim 10, characterized in
10 that the unit dose comprises from 100 to 1000 mg of metformin and from 50 to 600 mg of bezafibrate.



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/19, 31/215 // A61K 31:155	A3	(11) International Publication Number: WO 99/40904 (43) International Publication Date: 19 August 1999 (19.08.99)
(21) International Application Number: PCT/EP99/00614 (22) International Filing Date: 30 January 1999 (30.01.99) (30) Priority Data: 98/01709 12 February 1998 (12.02.98) FR (71) Applicant (for all designated States except US): MERCK PATENT GMBH [DE/DE]; Frankfurter Strasse 250, D-64293 Darmstadt (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): BONHOMME, Yves [FR/FR]; 21 avenue de la Paix, F-69260 Charbonnières les Bains (FR). BRIET, Philippe [FR/FR]; 216, avenue Félix Faure, F-69003 Lyon (FR). (74) Common Representative: MERCK PATENT GMBH; Post- fach, D-64271 Darmstadt (DE).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i> (88) Date of publication of the international search report: 30 September 1999 (30.09.99)
(54) Title: PHARMACEUTICAL COMPOSITION COMPRISING A COMBINATION OF METFORMIN AND FIBRATE, AND ITS USE FOR THE TREATMENT OF HYPERGLYCEMIA		
(57) Abstract The present invention relates to a pharmaceutical composition comprising, as active principles, (i) metformin optionally in the form of one of its pharmaceutically acceptable salts, and (ii) a fibrate chosen from fenofibrate and bezafibrate, in combination with one or more pharmaceutically acceptable excipients. These compositions are particularly suitable for the treatment of non-insulin-dependent diabetes.		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakistan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

BEST AVAILABLE COPY

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 99/00614

A. CLASSIFICATION OF SUBJECT MATTER IPC 6 A61K31/19 A61K31/215 //A61K31:155		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 6 A61K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	S. R. DE SILVA ET AL.: "Metformin and clofibrate in maturity onset diabetes mellitus: advantages of combined treatment." DIABET. METABOL., vol. 5, no. 3, 1979, pages 223-229, XP002083497 cited in the application	
A	FR 2 264 525 A (S.E.M.S.) 17 October 1975 (1975-10-17)	
A	EP 0 305 890 A (BOEHRINGER MANNHEIM) 8 March 1989 (1989-03-08)	
<input type="checkbox"/> Further documents are listed in the continuation of box C. <input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents : "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "Z" document member of the same patent family		
Date of the actual completion of the international search 13 August 1999		Date of mailing of the international search report 19/08/1999
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer Klaver, T

Form PCT/ISA/210 (second sheet) (July 1992)

BEST AVAILABLE COPY

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 99/00614

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
FR 2264525 A	17-10-1975	AT 346856 B	27-11-1978
		AT 217875 A	15-04-1978
		BE 826999 A	16-07-1975
		CA 1062280 A	11-09-1979
		CH 609669 A	15-03-1979
		DE 2511576 A	25-09-1975
		GB 1462316 A	26-01-1977
		JP 1034650 C	20-02-1981
		JP 50135043 A	25-10-1975
		JP 55026134 B	11-07-1980
		LU 72102 A	20-08-1975
		NL 7503515 A, B,	24-09-1975
		US 4080472 A	21-03-1978
EP 305890 A	08-03-1989	DE 3729209 A	09-03-1989
		AT 95694 T	15-10-1993
		AU 2158688 A	25-05-1989
		CA 1323575 A	26-10-1993
		DE 3884895 D	18-11-1993
		DK 472688 A	02-03-1989
		ES 2059457 T	16-11-1994
		IE 62189 B	28-12-1994
		JP 1071813 A	16-03-1989
		JP 2647918 B	27-08-1997
		PH 24568 A	03-08-1990
		PT 88383 A, B	31-07-1989
		US 5304575 A	19-04-1994

BEST AVAILABLE COPY